

# **COVID-19 Testing Specimen Submission Guidelines**

## **Specimen Collection**

Please review guidelines recommended by the Centers for Disease Control and Prevention to ensure the appropriate infection control precautions are in place **before** collecting any specimens (https://www.cdc.gov/coronavirus/2019-ncov/hcp/clinical-criteria.html).

Acceptable specimen types for collection:

### 1. Upper Respiratory Specimens

i. **Nasopharyngeal (NP) swab** only in viral transport media (VTM), Universal Transport Media (UTM) or Aptima Specimen Transfer Tube [Cat No. PRD-05110 or 301154C (green label)].

#### Please see the video for additional instructions on NP swab collection:

https://www.youtube.com/watch?v=DVJNWefmHjE

#### OR

ii. **Oropharyngeal (OP) swab** in VTM, UTM or Aptima Multitest Swab Specimen Collection Kit [Cat. No. PRD-03546 (orange label)]

#### OR

iii. **Nasal swab** in VTM, UTM or Aptima Multitest Swab Specimen Collection Kit [Cat. No. PRD-03546 (orange label)]

Swab specimens should be collected using only swabs with a synthetic tip, such as nylon or Dacron, and an aluminum or plastic shaft. Calcium alginate swabs are unacceptable and swabs with cotton tips and wooden shafts are not recommended. ESwabs in Amies media are currently <u>not</u> a valid specimen type for COVID-19 testing.

A health care facility that requires swabs or transport media should contact the DC DFS PHL via email (<u>DFS-COVID19@dc.gov</u>) to request collection supplies. Please indicate how many kits you require in the email. Swabs and transport media should only be used for testing conducted at DFS PHL.

### 2. Lower Respiratory Specimens

- i. *Sputum* Have the patient rinse the mouth with water and then expectorate deep cough directly into a sterile, leak-proof, screw-cap sputum collection cup or a sterile dry container.
- ii. *Bronchoalveolar lavage (BAL)* Collect 2-3 mL into a sterile, leak-proof, screw-cap sputum collection cup or sterile dry container.



## Specimen Storage

All specimens must be refrigerated (2-8°C) promptly after collection and couriered/shipped on cold packs within 72 hours. Specimens being held for >72 hours must be stored at -70°C and couriered/shipped on dry ice. Please do not freeze specimens if courier pick up will occur within 24 hours. It is very important that you contact <a href="mailto:DFS-COVID19@dc.gov">DFS-COVID19@dc.gov</a> as soon as the specimen is collected so that a courier can be ordered to pick up the specimen.

### **Paperwork**

Any specimen being sent through the DC DFS PHL must have the following paperwork accommodating the specimen:

- 1. DC DFS PHL External Chain of Custody (CoC)
- 2. DC DFS PHL Test Requisition Form

All paperwork can additionally be found on DC DFS PHL's website under forms and documents: https://dfs.dc.gov/publication/phl-forms-and-documents

Only one DC DFS PHL test requisition form (example provided below) and one chain of custody for each set of specimens is required for testing.

If you would like to be signed for the DFS PHL web portal which allow for electronic ordering and PDF report, please email <u>DFS-COVID19@dc.gov</u>

Please ensure that all specimens submitted and their respective test requisition form has the following information on it:

- Full name of patient
- Date of birth
- Unique patient identifier (e.g., medical record number, patient ID)
- Date and time of specimen collection

Incorrectly labeled requisitions and specimens will result in testing delays.

## Courier Request

Once specimens are ready for pick up and the appropriate paperwork is completed, please email <u>DFS-COVID19@dc.gov</u> using the following template to request a courier:

- Sample type (NP/OP)
- Storage conditions (refrigerated/frozen)
- Name of Facility
- Address of Facility (including the room number)
- Point of Contact
- Point of Contact Phone Number

Do NOT include patient identifiers in the body of the email.

Please contact the DC Department of Forensic Sciences Public Health Laboratory for questions pertaining to testing Phone: 202-727-8956 (Monday-Friday from 8:30am-5:30pm) | 202-868-6561 (after-hours calls) | Fax: 202-481-3936 Email: DFS-COVID19@dc.gov



# Example of PHL Test Requisition Form (1 per patient)



District of Columbia \* Department of Forensic Sciences \* Public Health Laboratory 401 E Street SW \* 4th Floor \* Washington, DC 20024 \* Phone (202) 727-8956 \* Fax (202) 481-3464 General Laboratory Services Request Form PHL Director: Anthony Tran, DrPH, MPH, D(ABMM) CLIA#: 09D0968273

*	$\star$	*

Patient Information  Last Name*			First Name*		Middle Initia	uired Informatio
Date of Birth*			Sex*		If Female, Pr	
(MM/DD/YYYY) Address			City*	le Female 0	ther Yes State*	No Unknow
Sample ID (Laboratory ID, Outbre	eak#, Zika#, etc.)*		Medic	al Record Number		
care provider with same						
umber requesting testing					NEW WIR	
Name of Submitting Hospital, Lab	boratory, or other Faci	lity*		Healthcare Provide	r NPI #*	
Health Care Provider	Last Name*			First Name*		
Address (include room)*	Requests will be fax Critical values will re		_	City*	State*	Zip*
Primary Contact	ast Name			First Name	•	•
(If not the Health Care Provider) Telephone #*	Secure Fax #**			Email		
(primary)	<u> </u>					
** Final report will be sent to the	fax number above					
Specimen Information						
Date of Collection* (MM/DD/YYY	(Y) Nasophar	ynx (NP) swab	and	e of Collection*	•	n, Bronchial Wa
Reason for Submission*	combined NP/0					eolar Lavage sp
Diagnostic		d in either UTM	or VTM	DC Health	Contact:	
Specimen Type (check all that app		. Dro		Turni I	Tyrna Deca	Seculta Careatica
☐ Blood Culture Bottle ☐ Is ☐ Blood Tube (Plasma, Serum or	colate Cary-Bla	ir E-Swab	, ப	wab UTM Other (spe	VTM Slide	Sterile Containe
Specimen Source*	nchoalveolar Lavage	Rronchial Wa	ash			Nasonhan/ny (N
Abscess Blood Broi Oropharynx (OP) NP/O	P □Plasma □ Reine □ Vaginal □ O	ectal Serum		Buccal CSF	Endocervical [	
Abscess Blood Broom Oropharynx (OP) NP/O	P Plasma Reine Vaginal 0	ectal Serum		Buccal CSF	Endocervical [	
Abscess Blood Broid Oropharynx (OP) NP/OID Section Sources approve for testing are highlighted	P Plasma Reine Vaginal O	ectal Serum		Buccal CSF	Endocervical Sputum, induced	
Abscess Blood Brown Oropharynx (OP) NP/O NSISSUE Wisseld Wisse	P Plasma Reine Vaginal O	ectal Serum Other (specify)	Spu	Buccal CSF tum, expectoration  MOLEC	Endocervical Sputum, induced	
Abscess Blood Brown Oropharynx (OP) NP/O NSISSUE Wisseld Wisse	P Plasma Reine Vaginal O	ectal Serum Other (specify)  Ebola (PCF	Spu R) <sup>+</sup> uenza (F	Buccal CSF tum, expectoration  MOLEC	Endocervical Sputum, induced	
Abscess Blood Broot Oropharynx (OP) NP/Ot	P Plasma Reine Vaginal O	Ebola (PCF	R) <sup>+</sup> uenza (F	Buccal CSF tum, expectoration  MOLEC	Endocervical Sputum, induced	
Abscess Blood Broot Oropharynx (OP) NP/OT Tissue Urestral Ures specimen sources approve for testing are highlighted Test Request (* requested to BT RULE-OUT T/O B. anthracis  r/O Brucella sp. r/O Burkholderia sp. r/O F. tularensis	P Plasma Reine Vaginal O	Etal Serum S	R) <sup>†</sup> uenza (F (PCR) st Respi	Buccal CSF tum, expectoration  MOLEC  PCR)*	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR)†	Stool Thro
Abscess Blood Broot Oropharynx (OP) NP/OT NP/OT NESURE Wisseld Wisseld NP/OT NP/OT NESURE NES	P Plasma Reine Vaginal O	Etal Serum S	R)+ uenza (FCR) st Respi	Buccal CSF tum, expectoration  MOLEC	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR)†	Stool Thro
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NSISSUE Wisseld Wisseld No. 1 NP/O NSISSUE NI NEST NEW	P Plasma Reine Vaginal O	Ebola (PCF   Novel Influence   Novel Influence   Novel Influence   Novel Influence   Chlamydid   Mumps (P	R) <sup>+</sup> uenza (F (PCR) st Respi	Buccal CSF tum, expectoration  MOLEC  PCR)*  ratory Syndrome (I matis and Neisseria	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR)†	Stool Thro
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NS NP/O	P Plasma Reine Vaginal O	Ebola (PCF Novel Influence Middle Ea Chlamydid Mumps (P Measles V	R) <sup>+</sup> uenza (F (PCR) st Respi a tracho CCR) <sup>+</sup> firus (PC	Buccal CSF tum, expectoration  MOLEC  PCR)*  ratory Syndrome (I matis and Neisseria	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR) <sup>+</sup> a gonorrhoeae (TM	Stool Thro
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NS NP/O NS NP/O NS NP/O NS NP/O NS	P Plasma Reine Vaginal O	Ebola (PCF Novel Influence Middle Ea Chlamydid Measles V Arbovirus	R)* uenza (F (PCR) st Respi r tracho PCR)* firus (PC Detecti	Buccal CSF tum, expectoration  MOLEC  PCR)*  ratory Syndrome (I matis and Neisseria	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR) <sup>+</sup> a gonorrhoeae (TM	Stool Thro
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NS NP/O	P Plasma Reine Vaginal O	Ebola (PCF Novel Influence Minimum Middle Ea Chlamydid Mumps (P Measles V Arbovirus COVID-19	R)* uenza (F (PCR) st Respi r tracho PCR)* firus (PC Detecti	Buccal CSF tum, expectoration  MOLEC  PCR)*  ratory Syndrome (I matis and Neisseria	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR) <sup>+</sup> a gonorrhoeae (TM	Stool Thro
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NS NP/O NS NP/O NS NP/O NS NP/O NS	P Plasma Reine Vaginal O	Ebola (PCF Novel Influence of Mumps (PCF) Middle Ea Chlamydid Mumps (PCF) Measles V Arbovirus COVID-19 SEROLOGY	Spu  R)* uenza (F (PCR) st Respi n tracho PCR)* firus (PC Detecti (PCR)*	Buccal CSF tum, expectoration  MOLEC  PCR)*  ratory Syndrome (I matis and Neisseria R)* on Panel (chikungu	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR) <sup>+</sup> a gonorrhoeae (TM	Stool Thro
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NS NP/O NS NP/O NS NP/O NS NP/O NS	P Plasma Reine Vaginal O	Ebola (PCF Novel Influence of Mumps (PCF) Middle Ea Chlamydid Mumps (PCF) Measles V Arbovirus COVID-19 SEROLOGY Measles V	R) <sup>+</sup> uenza (F (PCR) st Respira tracho PCR) <sup>+</sup> firus (PC Detecti (PCR) <sup>+</sup>	Buccal CSF tum, expectoration  MOLEC  PCR)*  ratory Syndrome (I matis and Neisseria R)* on Panel (chikungu	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR) <sup>+</sup> a gonorrhoeae (TM	Stool Thro
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NS NP/O NS NP/O NS NP/O NS NP/O NS	P Plasma Reine Vaginal O	Ebola (PCF Novel Influence of Mumps (PCF) Middle Ea Chlamydid Mumps (PCF) Measles V Arbovirus COVID-19 SEROLOGY	R) <sup>+</sup> uenza (F (PCR) st Respira tracho PCR) <sup>+</sup> firus (PC Detecti (PCR) <sup>+</sup>	Buccal CSF tum, expectoration  MOLEC  PCR)*  ratory Syndrome (I matis and Neisseria R)* on Panel (chikungu	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR) <sup>+</sup> a gonorrhoeae (TM	Stool Thro
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NS NP/O NS NP/O NS NP/O NS NP/O NS	P Plasma Reine Vaginal O	Ebola (PCF Novel Influence of Middle Ea Chlamydia Measles V Arbovirus COVID-19 SEROLOGY Measles V Zika Virus	R)+ uenza (F (PCR) r tracho PCR)+ l'irus (PC Detecti (PCR)+ l'irus (IgC (IgM)+	Buccal CSF tum, expectoration  MOLEC  CCR)*  ratory Syndrome (I matis and Neisseria R)* on Panel (chikungu	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR)+ a gonorrhoeae (TM  unya, dengue and 2	Stool Thro
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NS NP/O NS NP/O NS NP/O NS NP/O NS	P Plasma Reine Vaginal O	Ebola (PCF  Novel Influ Norovirus  Middle Ea Chlamydia Measles V Arbovirus COVID-19 SEROLOGY Measles V Zika Virus	R)+ uenza (F (PCR) r tracho PCR)+ firus (PC Detecti (PCR)+ firus (Ig( (IgM)+ ry DFA v	Buccal CSF tum, expectoration  MOLEC  PCR)*  ratory Syndrome (I matis and Neisseria R)* on Panel (chikungu	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR)+ a gonorrhoeae (TM  unya, dengue and 2  ULTURE Culture	IA)
Abscess Blood Broom Cropharynx (OP) NP/O NP/O NS NP/O NS NP/O NS NP/O NS NP/O NS	P Plasma Reine Vaginal O	Ebola (PCF  Novel Influ Norovirus  Middle Ea Chlamydia Measles V Arbovirus COVID-19 SEROLOGY Measles V Zika Virus	R)* uenza (F (PCR) st Respia tracho CCR)* firus (PC Detecti (PCR)* firus (IgC (IgM)* ry DFA v spiratory:	Buccal CSF tum, expectoration  MOLEC  PCR)*  ratory Syndrome (I matis and Neisseria R)* on Panel (chikungu	Endocervical Sputum, induced  ULAR  MERS-CoV) (PCR)+ a gonorrhoeae (TM  unya, dengue and 2  ULTURE Culture	IA)